arXiv:1504.00055v1 [math.DG] 31 Mar 2015

MEAN CURVATURE, VOLUME AND PROPERNESS OF ISOMETRIC IMMERSIONS

VICENT GIMENO* AND VICENTE PALMER**

ABSTRACT. We explore the relation among volume, curvature and properness of a m-dimensional isometric immersion in a Riemannian manifold. We show that, when the L^p-norm of the mean curvature vector is bounded for some $m \leq p \leq \infty$, and the ambient manifold is a Riemannian manifold with bounded geometry, properness is equivalent to the finiteness of the volume of extrinsic balls. We also relate the total absolute curvature of a surface isometrically immersed in a Riemannian manifold with its properness. Finally, we relate the curvature and the topology of a complete and non-compact 2-Riemannian manifold M with non-positive Gaussian curvature and finite topology, using the study of the focal points of the transverse Jacobi fields to a geodesic ray in M . In particular, we have explored the relation between the minimal focal distance of a geodesic ray and the total curvature of an end containing that geodesic ray.

1. INTRODUCTION

In 1965, E. Calabi stated in [5] the following conjecture, (also known as "the first Calabi's conjecture"): *prove that a complete minimal hypersurface in* $\mathbb{R}^n$ *must be unbounded*. This conjecture turned to be false when we consider *immersed* hypersurfaces: in the paper [28], N. Nadirashvili constructed a complete immersion of a minimal disk into the unit ball in $\mathbb{R}^3$. Nadirashvili's complete immersion is not proper, so it is natural to wonder about the rôle of properness in relation with the first Calabi's conjecture.

In fact, in the paper [11], T. H. Colding and W. P. Minicozzi studied the following question (also known as the Calabi-Yau conjecture): *Prove that every complete embedded minimal surface in* $\mathbb{R}^3$ *is proper.*

In [11] the authors proved that a complete embedded minimal disk in $\mathbb{R}^3$ must be proper, so the first Calabi's conjecture is true for embedded minimal disks. Moreover they proved that a complete embedded minimal surface with finite topology, (namely, homeomorphic to a compact 2-dimensional manifold, from which it has been removed a finite number of points, which corresponds to the number of *ends* of the surface), is properly immersed in $\mathbb{R}^3$.

On the other hand, W. H. Meeks and H. Rosenberg proved in the paper [25] that a complete embedded minimal surface with positive injectivity radius in $\mathbb{R}^3$ must be proper. This result includes Colding's-Minicozzi result because they proved too in this paper that every complete embedded minimal surface with finite topology in $\mathbb{R}^3$ has positive injectivity radius. We shall extract the intrinsic aspects of this proof (see Lemma 4.1) to show our Theorem 1.4.

More recently, Calabi-Yau's conjecture has been explored in broader contexts, namely, W. H. Meeks, J. Perez and A. Ros have proved, (see [26]), that Calabi-Yau conjecture is true when we consider embedded minimal surfaces with finite genus and countable number of ends in $\mathbb{R}^3$. On the other hand, B. Coskunuzer, W. H. Meeks and G. Tinaglia proved that Calabi-Yau conjecture is not true in $\mathbb{H}^3$: there exists a non properly embedded minimal

---

* Work partially supported by the Research Program of University Jaume I Project P1-1B2012-18, and DGI -MINECO grant (FEDER) MTM2013-48371-C2-2-P.

**Work partially supported by the Research Program of University Jaume I Project P1-1B2012-18, DGI -MINECO grant (FEDER) MTM2013-48371-C2-2-P, and Generalitat Valenciana Grant PrometeoII/2014/064 .

plane (genus zero, one end) in $\mathbb{H}^3$, (see [13]). In this way, M. Rodriguez and G. Tinaglia have constructed a family of non-proper complete minimal disks with finite total curvature in the quotient of $\mathbb{H}^2 \times \mathbb{R}$ by the discrete group of isometries generated by vertical translations and screw motions, which are embedded in $\mathbb{H}^2 \times \mathbb{R}$, (see the paper [30]).

Taking as a starting point all these results, several questions arises in a natural way, e.g., what happens when we consider non minimal surfaces (or submanifolds) immersed in a more general ambient space?. Or, could the embeddedness of the immersion be replaced by another hypothesis related with the volume or the metric?. Summarizing: what can be said about geometric conditions describing the properness of an immersion in a broader context?

Looking for these geometric conditions and given an isometric immersion $\varphi : P^m \to N^n$, where P^m and N^n are complete Riemannian manifolds, a quite natural assumption to be considered is following

$$\text{Vol}(\varphi^{-1}(B_t^N)) < \infty, \quad \forall t > 0 . \tag{1.1}$$

Where B_t^N is the geodesic t-ball in the ambient manifold N. The domains $D_t = \varphi^{-1}(B_t^N)$ are known as the *extrinsic* t-balls.

In fact, when the ambient manifold is complete, inequality (1.1) is a necessary condition for properness. However, it is not a sufficient condition: consider e. g. a negatively curved and non-compact surface with finite volume which can be immersed (via Nash immersion theorem) in a metric ball in $\mathbb{R}^N$ for large N.

On the other hand there are immediate examples of proper non-embedded minimal surfaces in $\mathbb{R}^3$, such as Enneper's surface or proper minimal surfaces with infinite topology, such the 1-periodic Scherk's surface which in its turn satisfies inequality (1.1).

We can find in the very recent literature, (see Remark 4 in [24]), the proof that inequality (1.1) satisfied by the extrinsic balls D_t of a minimal immersion $\varphi : P^m \to N^n$ in a Cartan-Hadamard manifold N^n with sectional curvatures bounded from above $K_N \leq b \leq 0$ implies the properness of P^m. An older reference of the same result, but for minimal surfaces in $\mathbb{R}^3$ with finite topology can be found in Theorem 3 in [9].

Concerning the techniques used, the proof in [24] is based in the intrinsic monotonicity property satisfied by the (intrinsic) volume growth $\frac{\text{Vol}(B_t^N)}{\text{Vol}(B_t^{b,m})}$ of such immersions. Here, we shall denote as $B_t^{b,m}$ the geodesic t-ball in the real space form $\mathbb{K}^m(b)$, and as $S_t^{b,m-1}$ the geodesic t-sphere in $\mathbb{K}^m(b)$. On the other hand, we can find also in Lemma 4 in [35] an extrinsic approach to the use of the monotonicity formula of the extrinsic volume growth $\frac{\text{Vol}(D_t)}{\text{Vol}(B_t^{b,m})}$ to prove that a minimal hypersurface $\varphi : P^{n-1} \to \mathbb{R}^n$ is proper when the extrinsic volume growth is finite, (a more restrictive hypothesis than our condition (1.1)).

This approach inspires the interplay between the intrinsic and the extrinsic distances used in [24] to prove the properness of a minimal immersion in a Cartan-Hadamard manifold as well as our own way to prove Theorem 1.1, where we obtain a characterization of properness for non-minimal immersions in a broader family of ambient Riemannian manifolds, namely, the Riemannian manifolds with *bounded geometry*.

Recall that a complete and non-compact Riemannian manifold N is a manifold with *bounded geometry* if its sectional curvatures K_N are bounded from above, $K_N \leq b$, ($b \in \mathbb{R}^+$), and its injectivity radius $i_0(N) = \text{Inf}\{i_N(p) : p \in N\}$ is positive $i_0(N) = i_0 > 0$. We must remark that Cartan-Hadamard manifolds are simply connected Riemannian manifolds with bounded geometry, being $b = 0$ and $i_0(N) = \infty$ in this case.

In fact, we have used a comparison for the volume of geodesic balls in a submanifold with controlled L^p norm of its mean curvature immersed in such ambient manifolds, stablished by M.P. Cavalcante, H. Mirandola and F. Vitorio in [7] as a part of the proof of their Theorem B. We follow a similar vein of reasoning that these authors, but for different purposes: while they prove that the volume of the ends of the submanifold is infinite, we characterize the properness of the immersion (see Remarks 1.2 and 3.2 where we try to relate both facts).

In this regard, we have obtained the following theorem for submanifolds with bounded L^p-norm of their mean curvature vector, immersed in a Riemannian manifold with bounded geometry.

Theorem 1.1. *Let $\varphi : P^m \longrightarrow N^n$ be an isometric immersion of a complete non-compact manifold P^m in a manifold N with bounded geometry. Assume that the mean curvature vector of φ satisfies $\|\vec{H}_P\|_{L^p(P)} < \infty$, for one $m \leq p \leq \infty$.*

Then, P^m is properly immersed in N^n if and only if $\mathrm{Vol}(\varphi^{-1}(B_t^N)) < \infty$ for all $t > 0$, where B_t^N denotes any geodesic ball of radius t in the ambient manifold N.

Remark 1.2. Applying Theorem B of [7] to an isometric immersion $\varphi : P^m \longrightarrow N^n$ as in Theorem 1.1 we have that $\mathrm{Vol}(P) = \infty$. In Remark 3.2 we shall extend this comment to the ends of P, as in Theorem B in [7].

As a corollary, we have immediately that CMC surfaces are properly immersed when inequality (1.1) is satisfied:

Corollary 1.3. *Let $\varphi : P^m \longrightarrow N^n$ be an isometric immersion of a complete non-compact manifold P^m in a manifold N with bounded geometry. Assume that the mean curvature vector of φ has constant length.*

Then, P^m is properly immersed in N^n if and only if, for all $t > 0$, the extrinsic ball $\varphi^{-1}(B_t^N)$ of radius t has finite volume $\mathrm{Vol}(\varphi^{-1}(B_t^N)) < \infty$.

A more classical approach to the study of the relationship between geometric invariants and properness of an immersion $\varphi : P^m \to N^n$ encompasses bounds on the total extrinsic curvature $\int_P \|B^P\|^2 d\sigma$, where B^P denotes the second fundamental form of the immersion. Note that when we consider minimal immersions of dimension two in the Euclidean space $\mathbb{R}^n$, this total extrinsic curvature agrees with total absolute curvature $\int_P |K_P| d\sigma$, where K_P denotes the Gaussian curvature of the surface P.

Concerning this relation among the total (extrinsic) curvature of an immersion and its properness, S. Muller and V. Sverak proved in [27] that a complete and oriented surface $\varphi : M^2 \to \mathbb{R}^n$ with finite total extrinsic curvature $\int_M \|B^M\|^2 d\sigma < \infty$ is proper.

Within this study of the behavior at infinity of complete and minimal immersions with finite total extrinsic curvature, it was also proven by M. T. Anderson in [1] and by G. de Oliveira in [29] that the immersion of a complete and minimal submanifold P^m in $\mathbb{R}^n$ or $\mathbb{H}^n(b)$ satisfying $\int_P \|B^P\|^m d\sigma < \infty$ is proper and that P^m has finite topology.

At this point we should mention the results by G. P. Bessa et al. reported in [2] and in [3], where new conditions have been stated on the decay of the extrinsic curvature for a completely immersed submanifold P^m in the Euclidean space ([2]) and in a Cartan-Hadamard manifold ([3]), which guarantees the properness of the submanifold and the finiteness of its topology.

In the second of our main results we deal with the relation among total curvature and properness, proving that finiteness of total curvature is not a necessary condition for properness. In order to state the theorem, we recall that given a compact subset $D \subset M$ of a Riemannian manifold M, an *end* E M with respect to D is a connected unbounded component of $M \setminus D$. From now on, when we say that E is an end, it is implicitly assumed that E is an end with respect to some compact subset $D \subset M$ (see [23] and [14]).

Theorem 1.4. *Let $\varphi : M^2 \longrightarrow N^n$ be a complete Riemannian 2-manifold immersed in a complete Riemannian manifold N^n. Suppose that M has finite topological type, and has non-positive Gaussian curvature, $K_M \leq 0$. Let us suppose moreover that for any $t > 0$,*

$$\mathrm{Vol}(\varphi^{-1}(B_t^N)) < \infty. \tag{1.2}$$

If every end E of M satisfies $\int_E |K_M| d\sigma = \infty$ then M^2 is properly immersed in N^n.

Remark 1.5. Observe that:

1. This theorem can be stated on an end $E \subseteq M$ of a complete Riemannian 2-manifold immersed in a complete Riemannian manifold N^n, $\varphi : M^2 \to N^n$, assuming that $K_M \leq 0$ on E, that $\operatorname{Vol}(\varphi^{-1}(B_t^N) \cap E) < \infty \; \forall t > 0$ and that $\int_E |K_M| d\sigma = \infty$, and concluding that E is properly immersed in N.
2. None of the three assumptions of this result, namely, finiteness of the volume of extrinsic balls, non-positive Gaussian curvature of the submanifold and infinite total curvature of its ends, can be avoided, as we can see in the following examples.

   In first place, we need to assume that the volume of the extrinsic balls $\varphi^{-1}(B_t^N)$ is finite for all radius $t > 0$. An example of non-proper immersion with non-positive curvature, infinite total curvature and having extrinsic balls with infinite volume is Nadirashvili's minimal labyrinth $\varphi : M^2 \longrightarrow \mathbb{R}^3$ (see [28]). This surface has positive first eigenvalue of the Laplacian, $\lambda_1(M) > 0$, so it is hyperbolic, (see [20]). Hence, it has infinite total curvature and infinite volume, (see [21] and [34]). On the other hand, we have that $\varphi(M) \subseteq B_1^{\mathbb{R}^3}$, so $\operatorname{Vol}(\varphi^{-1}(B_1^{\mathbb{R}^3})) = \operatorname{Vol}(M) = \infty$.

   Other example of surfaces which do not satisfy the hypothesis on the volume of the extrinsic balls in our theorem are the helicoidal-Scherk examples constructed in [30]. They are simply connected minimal surfaces $\varphi : M^2_{n,h} \to \mathbb{H}^2 \times \mathbb{R}$ embedded in $\mathbb{H}^2 \times \mathbb{R}$ which are not properly embedded. They are invariant under discrete subgroups of isometries G of $\mathbb{H}^2 \times \mathbb{R}$ generated by translations and screw motions, and if we consider the quotient of $M^2_{n,h}$ by a subgroup of G, these surfaces have finite total curvature. However, the surfaces $M^2_{n,h}$ have infinite total curvature. They also have non-positive Gaussian curvature. These surfaces accumulates to a subset of a cylinder in $\mathbb{H}^2 \times \mathbb{R}$, so we have, fixing a pole $o \in \mathbb{H}^2 \times \mathbb{R}$ and using the argument given in the proof of Theorem 1.1 (recall that $M_{n,h}$ is minimal), a sequence of points diverging in $M_{n,h}$ and converging to a limit point in $\mathbb{H}^2 \times \mathbb{R}$. We can consider now a sequence of (disjoint) geodesic balls centered in these points. It is easy to see that the union B of all these geodesic balls is included in some extrinsic ball $\varphi^{-1}(B_{R_0}^{\mathbb{H}^2 \times \mathbb{R}})$. We apply Lemma 3.1 to conclude that $\operatorname{Vol}(\varphi^{-1}(B_{R_0}^{\mathbb{H}^2 \times \mathbb{R}})) \geq \operatorname{Vol}(B) = \infty$.

   Secondly, all the ends of the submanifold must have non-finite total absolute curvature. Let us consider a complete and non-compact Riemannian manifold M with dimension two, with constant Gaussian curvature -1 and finite volume $\operatorname{Vol}(M) < \infty$. As we mentioned above, a surface like this can be isometrically immersed (via Nash immersion theorem) in a metric ball B_1^N in $\mathbb{R}^N$ for large N. Obviously, is not properly immersed, it has negative curvature -1 and if we consider any end $E \subseteq M$, we have that $\int_E |K_M| d\sigma \leq \int_E |K_M| d\sigma = \operatorname{Vol}(M) < \infty$.

   Finally, and concerning the assumption of non-positiveness of the Gaussian curvature, we can consider the computational example given by a modified pseudoesphere S^2, isometrically immersed in $\mathbb{R}^3$, given by the following parametrization

$$\begin{cases} x = 2\left(1 + \operatorname{sech}(u)\cos(v)\right) \cdot \cos(u - \tanh(u)) \\ y = 2\left(1 + \operatorname{sech}(u)\cos(v)\right) \cdot \sin(u - \tanh(u)) \\ z = 2\operatorname{sech}(u)\sin(v) \end{cases} \tag{1.3}$$

   for $(u, v) \in [0, \infty) \times [0, 2\pi)$. The two-dimensional volume of the extrinsic balls of S^2 is finite because, as can be showed numerically, the total two-dimensional volume $\operatorname{Vol}(S^2)$ of S^2 is finite, $\operatorname{Vol}(S^2) \approx 25.509$. Moreover, its total curvature is infinite. This immersion is not proper because S^2 accumulates to a circle (see



FIGURE 1. Modified pseudoesphere given by parametrization (1.3). In the left side $u \in [0, 10]$, and in the right $u \in [0, 15]$. This surface has finite total area, and hence, finite area of its extrinsic balls, and it has infinite total curvature, $\int_{S^2} |K| d\sigma = \infty$. But this immersion is not proper because the image accumulates in the circle of radius 2 in the plane $\{(x, y, z) \in \mathbb{R}^3 \,:\, z = 0\}$. This example shows that the hypothesis $K_M \leq 0$ is needed in order to state theorem 1.4.

figure 1), but we must notice that S^2 has points where its Gaussian curvature is positive.

To prove Theorem 1.4 we use Lemma 4.1, which is an intrinsic version of the proof, given in [25], that the injectivity radius $\mathrm{inj}(M)$ of a complete embedded minimal surface M in $\mathbb{R}^3$ of finite topology is positive. In this paper the authors proved that, if we assume that $\mathrm{inj}(M) = 0$, then there exists an end of the surface $E \subseteq M$ that has finite absolute total curvature.

A corollary of Theorem 1.4 and the fact that every 2-dimensional manifold with finite total absolute curvature is parabolic (see [21]) is following:

Corollary 1.6. *Let $\varphi : M^2 \longrightarrow N^n$ be a complete Riemannian 2-manifold immersed in a complete Riemannian manifold N^n. Suppose that M has finite topological type, non-positive Gaussian curvature, $K_M \leq 0$, and that all its ends are hyperbolic. Let us suppose moreover that for any $t > 0$,*

$$\mathrm{Vol}(\varphi^{-1}(B_t^N)) < \infty. \tag{1.4}$$

Then, M^2 is properly immersed in N^n.

We have adapted, on the other hand, the techniques used in Lemma 4.1, to relate the curvature and the topology of a complete and non-compact 2-Riemannian manifold, using of the notion of *geodesic ray* and the study of the focal points of the transverse Jacobi fields to a submanifold. In particular, we have explored the relation between the minimal focal distance of a geodesic ray and the total curvature of an end containing that geodesic ray.

Before to state our last main result, we present some definitions. Recall that, given a Riemannian manifold (M^n, g), a geodesic $\gamma : [0, \infty) \to M$ emanating from $p \in M$ parametrized by arc-length is called a *ray* emanating from p if

$$\mathrm{dist}_M(\gamma(t), \gamma(s)) = |t - s|$$

for all $t, s \geq 0$ (see for instance [32]). Whenever one has a geodesic ray γ in a surface M, one can construct, for any $t > 0$, the subset Ω_γ^t of the normal bundle $T\gamma^\perp$ of γ in M defined by

$$\Omega_\gamma^t := \left\{(p, \xi) \in T\gamma^\perp \,:\, |\xi| < t\right\}. \tag{1.5}$$

Denoting by $\exp^\perp$ the restriction of the exponential map to the normal bundle one can define the *minimal focal distance* $\mathrm{minfoc}(\gamma)$ of the geodesic ray γ as (see also §2 and

[17]) the maximum $t > 0$ such that the map

$$\exp^{\perp} : \Omega^{t}_{\gamma} \to \exp^{\perp}(\Omega^{t}_{\gamma}) \subset M \tag{1.6}$$

is a diffeomorphism.

Finally, we say that a geodesic ray γ emanating from $p \in M$ belongs to the end E (or equivalently, that E contains the ray γ), when $\gamma(t) \in E$ for any t large enough. At this point, we recall that an end can be also defined as an equivalent class of cofinal rays, (two rays in M, $\alpha(t)$ and $\beta(t)$ are *cofinal* iff for every compact set $K \subseteq M$, there exists t_0 such that if $t_1, t_2 \geq t_0$, then $\alpha(t_1), \beta(t_2)$ lie in the same connected component of $M - K$).

We have obtained the following intrinsic result:

Theorem 1.7. *Let M^2 be a complete, non compact and orientable surface with finite topology and non-positive Gaussian curvature, $K_M \leq 0$. Suppose that there exist a geodesic ray $\gamma \subseteq M$ with zero minimal focal distance,* $\operatorname{minfoc}(\gamma) = 0$. *Then* $\operatorname{inj}(M) = 0$ *and, moreover, γ belongs to an end $E_\gamma \subset M$ of finite total curvature, $\int_{E_\gamma} |K_M| d\sigma < \infty$.*

We shall finish this introduction with an example:

Example 1.8. When M has negative curvature, we can use [17, Corollary 8.6] to see that if $\operatorname{Vol}(M) < \infty$, then every ray γ has $\operatorname{minfoc}(\gamma) = 0$ (and hence by using theorem 1.7, every end of M has finite total curvature). However, there exist manifolds with negative curvature satisfying that every ray γ has $\operatorname{minfoc}(\gamma) = 0$ and with $\operatorname{Vol}(M) = \infty$, as it is easy to check considering the 2-Riemannian manifold $\Sigma = \mathbb{R} \times_w \mathbb{S}^1$ with warping metric $g_\Sigma = dt^2 + w^2(t)d\theta^2 = dt^2 + \sinh\left(-t + \sqrt{t^2+1}\right)^2 d\theta^2$.

Observe that Σ has finite topological type. Indeed, Σ has the topological type of $\mathbb{R} \times \mathbb{S}^1$ and, therefore Σ has zero genus and two ends. Given $\epsilon > 0$, let us consider the compact domain $\Omega_\epsilon \subset \Sigma$ given by $\Omega_\epsilon = [-\epsilon, \epsilon] \times \mathbb{S}^1$. Σ has, hence, two ends with respect to Ω_ϵ, namely:

$$\begin{aligned} E_- &:= (-\infty, -\epsilon) \times \mathbb{S}^1 \\ E_+ &:= (\epsilon, +\infty) \times \mathbb{S}^1. \end{aligned} \tag{1.7}$$

Both of the above ends are of infinite area $2\pi \int_\epsilon^\infty \sinh\left(-t + \sqrt{t^2+1}\right) dt = \infty$, and, since

$$\lim_{t \to \infty} \sinh\left(-t + \sqrt{t^2+1}\right) = 0, \tag{1.8}$$

the end E_+ is an shrinking end and hence every line in Σ has zero minimal focal distance. In fact, E_+ has finite total curvature, as it can be explicitly checked:

$$\begin{aligned} \int_{E_+} |K_\Sigma| dV &= \int_{E_+} \omega''(t) dt d\theta \\ &= 2\pi \int_\epsilon^\infty \Big((y'(t))^2 \sinh(y(t)) + y''(t)\cosh(y(t)) \Big) dt \approx 1.54308 \end{aligned} \tag{1.9}$$

when ϵ goes to zero.

1.1. **Outline.** In section §.2 we present some basic tools, based in the Jacobi index theory, we use: first, a key comparison result for the Laplacian of the extrinsic distance in a submanifold. Secondly, and to define the notion of *tube* around a submanifold immersed in a Riemannian manifold, we present the transverse Jacobi fields to a submanifold and the normal exponential map in relation of the minimal focal distance of this submanifold. In Section §.3 we state and prove Theorem 1.1. In Section §.4, we prove Theorem 1.4. Finally, we shall prove Theorem 1.7 in Section §.5. In Section §.6 we have studied some consequences of Theorem 1.7.

1.2. **Acknowledgments.** We would like to thank useful conversations with A. Alarcón, A. Hurtado, L. Jorge, F. López, and J. Pérez.

2. PRELIMINARIES

2.1. **Extrinsic distance balls.** Throughout the paper we assume that $\varphi : P^m \to N^n$ is an isometric immersion of a complete non-compact Riemannian m-manifold P^m into a complete Riemannian manifold N^n with bounded geometry. We shall refer P as a *submanifold* of N.

Given a point $o \in N$, for every $x \in N^n - Cut\{o\}$ we define $r_o(x) = \operatorname{dist}_N(o, x)$, and this distance is realized by the length of a unique geodesic from o to x, which is the *radial geodesic from o.*

When $\varphi^{-1}(N - Cut\{o\}) \neq \emptyset$, we also denote by r the composition

$$r_o \circ \varphi : \varphi^{-1}(N - Cut\{o\}) \subseteq P \to \mathbb{R}_+ \cup \{0\}$$

This composition is called the *extrinsic distance function* from o in P^m. The gradients of r_o in N and r in P are denoted by $\nabla^N r_o$ and $\nabla^P r$, respectively. Then we have the following basic relation, by virtue of the identification, given any point $x \in P$, between the tangent vectors $X \in T_xP$ and $\varphi_{*_x}(X) \in T_{\varphi(x)}N$

$$\nabla^N r_o = \nabla^P r + (\nabla^N r_o)^\perp \tag{2.1}$$

where $(\nabla^N r_o)^\perp(\varphi(x))$ is perpendicular to T_xP for all $x \in P$.

Definition 2.1. Given $\varphi : P^m \longrightarrow N^n$ an isometric immersion of a complete and connected Riemannian m-manifold P^m into a complete Riemannian manifold N^n with bounded geometry, we denote the *extrinsic metric balls* of radius $t > 0$, ($t \leq min\{i_0(N), \frac{\pi}{\sqrt{b}}\}$) and center $o \in N$ by $D_t(o)$. They are defined as the subset of P:

$$D_t(o) := \Big\{x \in P \,:\, r(x) < t\Big\} = \varphi^{-1}\left(B^N_t(o)\right)$$

where $B^N_t(o)$ denotes the open geodesic ball of radius t centered at the point o in N^n. Note that the set $\varphi^{-1}(o)$ can be the empty set.

The extrinsic balls are the sublevel sets of the extrinsic distance function.

2.2. **Transverse Jacobi fields and the normal exponential map to a submanifold.** Let $\varphi : P^m \to N^n$ be an isometric embedding of a complete non-compact Riemannian m-manifold P^m in a Riemannian manifold N. Let us identify P with the zero section $O(TP^\perp)$ of the normal bundle $TP^\perp$ of P in N. Then, (cfr. [32], [17]), we have that $\exp_{{TP^\perp}_{*(p,0)}} : T_{(p,0)}TP^\perp \to T_pN$ is an isomorphism for each $p \in P$, because for all $p \in P$, we have the decomposition $T_pN = T_pP \oplus T_pP^\perp = T_{(p,0)}TP^\perp$. Then, applying the inverse mapping theorem, we have that $\exp_{TP^\perp} : TP^\perp \to N$ is a diffeomorphism if it is restricted to an open neighborhood Ω_P of P in $TP^\perp$. We also shall denote $\exp_{TP^\perp}$ as $\exp^\perp$. Note that for any point $q \in \exp_{TP^\perp}(\Omega_P)$ there exists a unique minimal geodesic γ parametrized by arc length from a point of P to q which realizes the distance $\operatorname{dist}_N(q, P)$. This geodesic γ is perpendicular to P.

Definition 2.2. (See [17]). Let $\gamma_\xi(t)$ in N a geodesic normal to P at $t = 0$. The point $q = \gamma_\xi(t_0)$ is a *cut-focal point* along γ_ξ provided the distance from q to P is no longer minimized along γ_ξ after q. In other words, beyond the point q, there is a point in the geodesic $\tilde{q} \in \gamma_\xi$ such that the distance of $\tilde{q}$ to P is less than the distance of q to P, and it is realized by another geodesic $\tilde{\gamma}_\xi$ that contains $\tilde{q}$ and meets P orthogonally.

Definition 2.3. Given the unitary normal bundle $\mathbb{S}TP^\perp$ to P in N, *i.e.*,

$$\mathbb{S}TP^\perp := \left\{(p, \xi) \in TP^\perp \,:\, \|\xi\| = 1\right\}.$$

One has the following function $e_c : \mathbb{S}TP^\perp \to \mathbb{R}$ defined as

$$e_c(p, \xi) := \sup_{t>0} \left\{t \,:\, \operatorname{dist}_N(\exp^\perp(p, t\xi), P) = t\right\}.$$

Remark 2.4 (See [17]). The set Ω_P where $\exp^{\perp}$ is a diffeomorphism can be written as

$$\Omega_P = \left\{(p, t\xi) \in \mathbb{S}TP^{\perp}\ :\ 0 \leq t < e_c(p,\xi)\right\}.$$

Definition 2.5 (See [17]). The *minimal focal* distance of P in N is defined as

$$\operatorname{minfoc}(P) := \inf_{(p,\xi)\in \mathbb{S}TP^{\perp}} \{e_c(p,\xi)\}.$$

Remark 2.6. When P is a point p, the minimal focal distance coincides with the injectivity radius at p.

Definition 2.7. Given $\varphi : P^m \longrightarrow N^n$ a submanifold of a Riemannian manifold N, and a number $t \leq \operatorname{minfoc}(P)$. The *tube of radius* t about P in N is the set

$$T_t(P) := \{x \in N\ :\ \operatorname{dist}_N(x,P) \leq t\}.$$

Remark 2.8. Since $t \leq \operatorname{minfoc}(P)$ in the above definition and by virtue of Remark 2.4,

$$T_t(P) = \{x \in N\ :\ \exists \text{ a geodesic } \gamma \text{ starting at } x, \text{ length } L(\gamma) \leq t \text{ meeting } P \text{ orthogonally}\}$$

and moreover,

$$T_t(P) = \exp_{TP^{\perp}}\Big(\left\{(p,\xi) \in TP^{\perp}, \text{ with } \|\xi\| \leq t\right\}\Big) \subseteq \exp_{TP^{\perp}}(\Omega_P).$$

Definition 2.9 (See [32, 17]). Let $\gamma_\xi(t)$ be a geodesic in N normal to P at $t = 0$. The point $q = \gamma_\xi(t_0)$ is a *focal point* along γ_ξ if there exists a nonzero P-Jacobi field $Y(t)$ along γ_ξ such that $Y(t_0) = 0$. We call t_0 its *focal value*. Given $p \in P$ and $\xi \in \mathbb{S}TP^{\perp}$, let us denote as $e_f(p,\xi)$ the first of these focal values along the normal geodesic starting at p.

Remark 2.10 (See [32, Lemma 4.8 in Chapter II]). The point $q = \gamma_\xi(t_0)$ is a focal point along γ_ξ if and only if rank $\exp^{\perp}_{*(t_0\xi)} < m$. Then,

$$e_f(p,\xi) = \inf_{t>0}\left\{t\ :\ \text{rank}\left(\exp^{\perp}_{*(t\xi)}\right) < m\right\}.$$

Remark 2.11 (See [32, Lemma 2.11 in Chapter III]). Given $(p,\xi) \in \mathbb{S}TP^{\perp}$, the relation among focal points and cut-focal points along the geodesic normal to P, $\gamma_\xi(t)$, is given by:

$$e_c(p,\xi) \leq e_f(p,\xi) \tag{2.2}$$

The following Proposition shows that, when P is totally geodesic in N, and N has non-positive sectional curvatures, the normal exponential map is a local diffeomorphism, as occurs with the exponential map.

Proposition 2.12. *There are not focal points along a normal geodesic to a totally geodesic submanifold P in a manifold of non-positive curvature N.*

Proof. The proof is based on the convexity of the norm of Jacobi fields. Let $\gamma_\xi(t)$ be a geodesic in N normal to P at $t = 0$ (namely, $\gamma_\xi(0) = p \in P$ and $\gamma'_\xi(0) = \xi \in T_pP^{\perp}$). Let us suppose that $q = \gamma_\xi(t_0)$ is a focal point along γ_ξ, *i.e.*, there exist a nonzero P-Jacobi field $Y(t)$ along γ_ξ such that $Y(t_0) = 0$. Then, $Y(0) \in T_pP$ and $Y'(0) - A_\xi(Y(0)) \in T_pP^{\perp}$, being A_ξ the Weingarten map of P.
Let us define $F(t) = \langle Y(t), Y(t)\rangle$. Then, $F'(t) = 2\langle Y'(t), Y(t)\rangle$ and

$$\begin{aligned} F''(t) =& 2\|Y'(t)\|^2 + 2\langle Y''(t), Y(t)\rangle \\ =& 2\|Y'(t)\|^2 - 2K\left(\gamma'_\xi(t), Y(t)\right)\,\|\gamma'_\xi(t) \wedge Y(t)\|^2 \\ \geq& 0, \end{aligned} \tag{2.3}$$

because $K\left(\gamma'_\xi(t), Y(t)\right) \leq 0\ \forall t$ and

$$\begin{aligned} \langle Y''(t), Y(t)\rangle &= -\Big\langle R\left(Y(t), \gamma'_\xi(t)\right)\gamma'_\xi(t), Y(t)\Big\rangle \\ &= -K\left(\gamma'_\xi(t), Y(t)\right)\cdot\left\|\gamma'_\xi(t) \wedge Y(t)\right\|^2. \end{aligned} \tag{2.4}$$

On the other hand, $F(0) = \|Y(0)\|^2 \geq 0$ and $F(t_0) = \|Y(t_0)\|^2 = 0$. We are going to compute $F'(0)$. Since $Y'(0) - A_\xi(Y(0)) = Z(p) \in T_pP^\perp$ and $Y(0) \in T_pP$,

$$\begin{aligned} F'(0) =& 2\left\langle Y'(0)\,, Y(0)\right\rangle = 2\left\langle A_\xi\left(Y(0)\right),\, Y(0)\right\rangle \\ =& 2\left\langle B^P\left(Y(0), Y(0)\right),\, \xi\right\rangle \end{aligned} \tag{2.5}$$

But P is totally geodesic, so $F'(0) = 0$. Moreover, using inequality (2.3) and since $Y'(0) \neq 0$ (because $Y(t)$ is a nonzero vector field, see [15, Chap. 5, Corollary 2.5 and Remark 2.6]) then $F''(0) > 0$.

Hence, for $t > 0$ sufficiently small,

$$F(t) = F(0) + F'(0)t + \frac{F''(0)}{2}t^2 + O(t^3) > F(0) \geq 0$$

Since $F''(t) \geq 0$, then $F'(t_2) \geq F'(t_1)$ when $t_2 \geq t_1$. Therefore $F(t) > 0$ for all $t > 0$. □

2.3. **The volume of a tube around a curve.** Let us consider a curve $\sigma \subseteq N$ as a submanifold of the ambient Riemnnian manifold N, using the inclusion map $i : \sigma \longrightarrow N$. We have the following result, which is proved in [17],

Theorem 2.13 (See [17, Corollary 8.6]). *Let σ be a curve with finite length $L(\sigma)$ in a Riemannian manifold M^n. Then, if $K_M \leq 0$, we have, for $0 \leq r < \operatorname{minfoc}(\sigma)$, that*

$$\operatorname{Vol}(T_r(\sigma)) \geq \frac{(\pi r^2)^{\frac{n-1}{2}}}{(\frac{1}{2}(n-1))!}L(\sigma). \tag{2.6}$$

Remark 2.14. Note that $\frac{(\pi r^2)^{\frac{n-1}{2}}}{(\frac{1}{2}(n-1))!}L(\sigma)$ corresponds to the volume of the tube of radius r around a curve with length $L(\sigma)$ in $\mathbb{R}^n$.

Corollary 2.15. *Let γ be a geodesic ray on a complete manifold M with non-positive curvature $K_M \leq 0$. Suppose that $\operatorname{minfoc}(\gamma) > 0$, then for any $0 < r < \operatorname{minfoc}(\gamma)$,*

$$\operatorname{Vol}(T_r(\gamma)) = \infty.$$

Proof. Let us consider a partition of $\gamma = \cup_{i=1}^\infty \gamma_i$, where γ_i is a geodesic segment with finite length $L(\gamma_i)$. As $K_M \leq 0$, one has that, as a consequence of Hotelling's tube formula, (see also [17, Corollary 8.6] and the above theorem), that for any $0 < r < \operatorname{minfoc}(\gamma)$,

$$\operatorname{Vol}(T_r(\gamma_i)) \geq \frac{(\pi r^2)^{\frac{n-1}{2}}}{(\frac{1}{2}(n-1))!}L(\gamma_i) \;\; \forall i = 1, ..., \infty$$

Then, since $T_r(\gamma) = \cup_{i=1}^\infty T_r(\gamma_i)$, we have

$$\operatorname{Vol}(T_r(\gamma)) = \sum_{i=1}^\infty \operatorname{Vol}(T_r(\gamma_i)) \geq \frac{(\pi r^2)^{\frac{n-1}{2}}}{(\frac{1}{2}(n-1))!}\sum_{i=1}^\infty L(\gamma_i) = \infty \tag{2.7}$$

□

3. PROOF OF THEOREM 1.1

In this section we shall prove the first of the main results of this paper.

In order to help the reader, we shall present again the statement of the result:

Theorem. *Let $\varphi : P^m \longrightarrow N^n$ be an isometric immersion of a complete non-compact manifold P^m in a manifold N with bounded geometry. Assume that the mean curvature vector of φ satisfies $\|\vec{H}_P\|_{L^p(P)} < \infty$, for one $m \leq p \leq \infty$.*

Then, P^m is properly immersed in N^n if and only if $\operatorname{Vol}(\varphi^{-1}(B_t^N)) < \infty$ for all $t > 0$, where B_t^N denotes any geodesic ball of radius t in the ambient manifold N.

Proof. First of all we shall need the following estimate for the volume of the geodesic balls of a submanifold with bounded mean curvature immersed in a Riemannian manifold with bounded geometry given in the proof of Theorem B in the paper [7]:

Lemma 3.1 (See Theorem B in [7])**.** *Let $\varphi : P^m \longrightarrow N^n$ be an isometric immersion of a complete non-compact manifold P^m in a manifold N with bounded geometry. Assume that the mean curvature vector of φ satisfies $\|\vec{H}_P\|_{L^p(P)} < \infty$, for one $m \leq p \leq \infty$.*

Then, given a fixed $p_0 \in P$ there exists ρ_0 sufficiently large and a positive constant $K > 0$ such that, for all $p \in P - B^P_{2\rho_0}(p_0)$, and for all $\rho \leq \rho_0$, we have

$$\mathrm{Vol}(B^P_\rho(p)) \geq \frac{1}{2Km}\rho^m. \tag{3.1}$$

Now, to prove the result, let us suppose first that the immersion is proper. As N is complete, then $\varphi^{-1}(B^N_t)$ is compact for all $t > 0$ and hence, $\mathrm{Vol}(\varphi^{-1}(B^N_t)) < \infty$.

On the other hand, if we suppose that the immersion is not proper, then there exists a point $o \in N$ and geodesic R_0- ball in N, $\bar{B}^N_{R_0}(o)$, with compact closure, such that $\varphi^{-1}(\bar{B}^N_{R_0}(o))$ is not compact. As it is closed, then $\varphi^{-1}(\bar{B}^N_{R_0}(o))$ is not bounded in P. Therefore, let us consider $\{p_i\}_{i=1}^\infty$ a sequence on points in $\varphi^{-1}(\bar{B}^N_{R_0}(o))$, which diverges in P and such that $\{\varphi(p_i)\}_{i=1}^\infty$ converges to a limit point $p \in N$.

Given p_0 the first point of the sequence, and applying Lemma 3.1, there exists ρ_0 sufficiently large and a positive constant $K > 0$ such that, for all $p \in P - B^P_{2\rho_0}(p_0)$, and for all $\rho \leq \rho_0$, we have

$$\mathrm{Vol}(B^P_\rho(p)) \geq \frac{1}{2Km}\rho^m. \tag{3.2}$$

On the other hand, as the sequence $\{p_i\}_{i=1}^\infty$ diverges in P, we know that $\{p_i\}_{i=n_0}^\infty \subseteq P - B^P_{2\rho_0}(p_0)$ for some $n_0 \in \mathbb{N}$.

Let us consider now, given the quantity ρ_0, a subsequence of $\{p_n\}_{n=1}^\infty$ formed by points p_n satisfying $\mathrm{dist}_P(p_n, p_{n+1}) > \rho_0$ $\forall n \in \mathbb{N}$, (see Remark 3.2). Now, we shall consider the geodesic balls $B^P_{\rho_0/2}(p_n)$.

With these radius, the balls $B^P_{\rho_0/2}(p_n)$ are pairwise disjoint. Moreover, we have that, by Lemma 3.1, $\mathrm{Vol}(B^P_{\rho_0/2}(p_n)) \geq \frac{1}{2Km}(\frac{\rho_0}{2})^m$ for all $n \geq n_0$.

Let us see that $\cup_{n=n_0}^\infty B^P_{\rho_0/2}(p_n) \subseteq D_{R_0+\rho_0/2}(o) = \varphi^{-1}(B^N_{R_0+\rho_0/2}(o))$. For that, let us consider $x \in \cup_{n=n_0}^\infty B^P_{\rho_0/2}(p_n)$. Then, there exists $n_1 \geq n_0$ such that $x \in B^P_{\rho_0/2}(p_{n_1})$, so, as $p_{n_1} \in \varphi^{-1}(\bar{B}^N_{R_0}(o))$, we have

$$\begin{aligned}
\mathrm{dist}_N\,(\varphi(x), o) \leq &\,\mathrm{dist}_N\,(\varphi(x), \varphi(p_{n_1})) + \mathrm{dist}_N(\varphi(p_{n_1}), o)\\
\leq &\,\mathrm{dist}_P\,(x, p_{n_1}) + R_0\\
\leq &\,\rho_0/2 + R_0
\end{aligned} \tag{3.3}$$

Then, as we know, for all $n \geq n_0$, that $\mathrm{Vol}\left(B^P_{\rho_0/2}(p_n)\right) \geq \frac{1}{2Km}(\frac{\rho_0}{2})^m > 0$, then

$$\begin{aligned}
\mathrm{Vol}(D_{R_0+\rho_0/2}) \geq &\,\mathrm{Vol}\left(\bigcup_{n=n_0}^{\infty} B^P_{\rho_0/2}(p_n)\right)\\
\geq &\sum_{n=n_0}^{\infty} \mathrm{Vol}\left(B^P_{\rho_0/2}(p_n)\right) \geq \sum_{n=n_0}^{\infty} \frac{1}{2Km}(\frac{\rho_0}{2})^m = \infty
\end{aligned} \tag{3.4}$$

which is a contradiction with the hypothesis. □

Remark 3.2. When we consider $\{q_i\}_{i=1}^\infty$ a sequence on points which diverges in P, (in the sense that $\lim_{n\to\infty} d_P(q_1, q_n) = \infty$), and given a fixed point $p_0 \in P$ and the radius ρ_0 as in Lemma 3.1 it is always possible to choose a subsequence $\{q_k\}_{k=1}^\infty \subseteq P$ and one radius $\delta = d_P(q_1, q_2)$ such that $B^P_\delta(q_k) \cap B^P_\delta(q_{k'}) = \emptyset$ for all sufficiently large $k \neq k'$ and such that, for all $k \geq k_0$, $B^P_\delta(q_k) \subseteq P - B^P_{2\rho_0}(p_0)$ in such a way that

$$\mathrm{Vol}(P) \geq \mathrm{Vol}\left(\bigcup_{k=k_0}^{\infty} B^P_\delta(q_k)\right) \geq \sum_{k=k_0}^{\infty} \mathrm{Vol}\left(B^P_\delta(q_k)\right) \geq \sum_{k=k_0}^{\infty} \frac{1}{2Km}\delta^m = \infty$$

In particular, given a point $p_0 \in P$ and the radius ρ_0 as in Lemma 3.1, if we consider, as in the paper [7], an end $E \subseteq P$ with respect the ball $B^P_{2\rho_0}(p_0)$, we can choose a sequence of points $q_2, q_3, ...$ in P such that

$$q_k \in E \cap (B^P_{2k\rho_0}(p_0) - B^P_{(2k-1)\rho_0}(p_0))$$

Then, $\lim_{k\to\infty} d_P(q_2, q_k) = \infty$. Moreover, for all $0 < R \leq \rho_0$ and for all $k \geq 2$, $B^P_R(q_k) \subseteq E - B^P_{2k\rho_0}(p_0))$, because $d_P(q_k, p_0) > (2k-1)\rho_0 > 2\rho_0$, and $B^P_R(q_k) \cap B^P_R(q_{k'}) = \emptyset$. Hence, fixing a radius $0 < R \leq \rho_0$,

$$\begin{aligned}\mathrm{Vol}(E) \geq \mathrm{Vol}(E - B^P_{2\rho_0}(p_0)) &\geq \mathrm{Vol}\left(\bigcup_{k=2}^{\infty} B^P_R(q_k)\right) \\ &\geq \sum_{k=2}^{\infty} \mathrm{Vol}\left(B^P_R(q_k)\right) \geq \sum_{k=2}^{\infty} \frac{1}{2Km} R^m = \infty\end{aligned}$$

4. PROOF OF THEOREM 1.4

In this section we shall prove Theorem 1.4. In order to help the reader, and as in Section §.3, we shall present again the statement of the results.

4.1. A previous lemma: on the positiveness of the injectivity radius.

In the paper [25] it is proved that the injectivity radius $\mathrm{inj}(M)$ of a complete embedded minimal surface M in $\mathbb{R}^3$ of finite topology is positive, by showing that, if we assume that $\mathrm{inj}(M) = 0$, then there exists an end of the surface $E \subseteq M$ that has finite absolute total curvature so as E is complete and embedded, it is asymptotic to a half catenoid or the end of a plane and the injectivity radius of M restricted to E is bounded away from zero, which is a contradiction and hence $\mathrm{inj}(M) > 0$. From an intrinsic point of view, we can state this result as follows proving it with exactly the same argument that in [25]:

Lemma 4.1. *(See* [25]*) Let M be a complete and non-compact 2-Riemannian manifold with finite topology, non-positive curvature $K_M \leq 0$ and with zero injectivity radius $\mathrm{inj}(M) = 0$. Then, there exists and end $E \subseteq M$ such that $\int_E |K_M| d\sigma < \infty$.*

Proof. Recall that the injectivity radius of $p \in M$ is defined as the supremum

$$\mathrm{inj}_M(p) := \sup\left\{t > 0 \,|\, \exp_p : B^{T_pM}_t(0_p) \to B^M_t(p) \text{ is a diffeomorphism}\right\}$$

or, equivalently, $\mathrm{inj}_M(p) = \mathrm{dist}_M(p, C_p)$, being C_p the cut-locus of p in M. The injectivity radius of M is defined as $\mathrm{inj}(M) := \inf_{p\in M}\{\mathrm{inj}_M(p)\}$.

If we assume that $\mathrm{inj}(M) = 0$, there exists therefore a divergent sequence of points $\{p_n\}^\infty_{i=1}$ of M such that $\lim_{n\to\infty} \mathrm{inj}_M(p_n) = 0$. Because on the contrary, if $\lim_{n\to\infty} p_n = p_0 \in M$, since $\mathrm{inj}_M : M \to \mathbb{R}_+$ defined as $p \to \mathrm{inj}_M(p)$ is continuous, then

$$\mathrm{inj}_M(p_0) = \mathrm{inj}_M(\lim_{n\to\infty} p_n) = \lim_{n\to\infty} \mathrm{inj}_M(p_n) = 0,$$

and this is in contradiction with the fact that, for all $p \in M$, there exists a small positive radius $\epsilon_p > 0$ such that the exponential map $\exp_p|_{B^{T_pM}_{\epsilon_p}(0_p)}$ is a diffeomorphism onto an open set in M.

On the other hand, M^2 has finite topological type which is equivalent to be homeomorphic to the interior of a compact surface with boundary. Each connected component of this boundary is identified with an end of M and is homeomorphic in its turn to a punctured disk. We call these ends *annular* ends.

Hence, there exists an annular end E of M such that a divergent subsequence of $\{p_n\}_{n=1}^{\infty}$ is included in E.

Let γ_n be an embedded geodesic loop based at the point p_n, which is smooth except perhaps at p_n. This loop exist since M has non-positive curvature and hence there are not conjugate points aloneg γ. We know that the length of this loop is $2\,\mathrm{inj}_M(p_n)$ and that the external signed angle corresponding to the vertex p_n is $\theta_n \in (-\pi, \pi)$.

Now, we are going to see that γ_n is not the boundary of a disk in M. If γ_n would be the boundary of a disk in M, then Gauss-Bonnet theorem should imply that

$$\int_{\Omega} K_M d\sigma = 2\pi\chi(\Omega) - \theta_n > 0, \tag{4.1}$$

where we have used that $\chi(\Omega) = 1$ because Ω is assumed to be a disk. But taking into account that $K_M \leq 0$, the above inequality is a contradiction.

Now, let us consider a compact annulus $E(n) \subset E$ bounded by the geodesic loops T_1 and T_n. Applying global Gauss-Bonnet theorem to the domain $E(n)$, we have

$$0 = 2\pi\chi\left(E(n)\right) = \int_{E(n)} K_M d\sigma + \theta_1 + \theta_n < \int_{E(n)} K_M d\sigma + 4\pi \tag{4.2}$$

Since the above inequality does not depend on n, E has finite absolute total curvature and the lemma follows. □

Remark 4.2. Observe that from the proof of the above lemma if we have a divergent sequence $\{p_i\}_{i=1}^{\infty}$ of points contained in an annular end $E \subset M$, of a surface M with non-positive curvature, and the sequence is such that

$$\lim_{n\to\infty} \mathrm{inj}(p_n) = 0.$$

Then

$$\int_E |K_M|\, d\sigma < \infty.$$

4.2. **Proof of Theorem 1.4.** We are going to prove Theorem 1.4, which asserts:

Theorem. *Let $\varphi : M^2 \longrightarrow N^n$ be a complete Riemannian 2-manifold immersed in a complete Riemannian manifold N^n. Suppose that M has finite topological type, and has non-positive Gaussian curvature, $K_M \leq 0$. Let us suppose moreover that for any $t > 0$,*

$$\mathrm{Vol}(\varphi^{-1}(B_t^N)) < \infty. \tag{4.3}$$

If every end E of M satisfies $\int_E |K_M| d\sigma = \infty$ then M^2 is properly immersed in N^n.

Proof. Let us suppose that M is not properly immersed. Then, there exists a geodesic ball in N, $\bar{B}_{R_0}^N(o)$, with compact closure, such that $\varphi^{-1}(\bar{B}_{R_0}^N(o))$ is not compact. Since it is closed, then $\varphi^{-1}(\bar{B}_{R_0}^N(o))$ is not bounded in M.

Therefore, as in the proof of Theorem 1.1, let us consider $\{p_i\}_{i=1}^{\infty} \subseteq \varphi^{-1}(\bar{B}_{R_0}^N(o))$ a sequence on points which diverges in M and such that $\{\varphi(p_i)\}_{i=1}^{\infty}$ converges to a limit point $p \in N$. Since the topology of M is finite, then there exists an annular end E such that a divergent subsequence of $\{p_n\}_{n=1}^{\infty}$ is included in E. Denoting as $\{q_n\}_{n=1}^{\infty}$ such subsequence, and given a positive quantity ϵ_1, let us choose a subsequence of $\{q_n\}_{n=1}^{\infty}$ formed by points q_n satisfying $\mathrm{dist}_M(q_n, q_{n+1}) > \epsilon_1$.

If every end E has infinite total curvature, then applying Lemma 4.1, we have that $\mathrm{inj}(M) = \delta > 0$. Hence, given the sequence $\{q_n\}_{n=1}^{\infty}$ formed by points q_n satisfying $\mathrm{dist}_M(q_n, q_{n+1}) > \epsilon_1$, we have that $\inf\{\mathrm{inj}_M(q_i)\}_{i=1}^{\infty} \geq \delta > 0$, so we shall consider now the geodesic balls $B_\epsilon^M(q_i)$ with $\epsilon < \min\{\delta, \frac{\epsilon_1}{2}\}$, in such a way that they are pairwise disjoint.

Now, we proceed as in the proof of Theorem 1.1 again, showing in the same way that

$$\bigcup_{i=1}^{\infty} B_\epsilon^M(q_i) \subseteq D_{R_0+\epsilon} = \varphi^{-1}(B^N_{R_0+\epsilon}(o))$$

Then, since $K_M \leq 0$ and $\epsilon < \text{inj}\,(M)$, we have, by using Bishop's-Gromov comparison theorem, that $\text{Vol}(B_\epsilon^M(q_i)) \geq \text{Vol}(B_\epsilon^{\mathbb{R}^n})$, thus

$$\begin{aligned} \text{Vol}\left(\varphi^{-1}(B^N_{R_0+\epsilon}(o))\right) &\geq \text{Vol}\left(\bigcup_{i=1}^{\infty} B_\epsilon^M(q_i)\right) \\ &\geq \sum_{i=1}^{\infty} \text{Vol}\left(B_\epsilon^M(q_i)\right) \geq \sum_{i=1}^{\infty} \text{Vol}\left(B_\epsilon^{\mathbb{R}^n}\right) = \infty \end{aligned} \tag{4.4}$$

which is a contradiction with the hypothesis. Hence, M is properly immersed.

□

5. PROOF OF THEOREM 1.7

In this section, we shall prove Theorem 1.7, which asserts:

Theorem. *Let M be a complete, non compact and orientable surface with finite topology and non-positive Gaussian curvature, $K_M \leq 0$. Suppose that there exist a geodesic ray $\gamma \subseteq M$ with zero minimal focal distance,* $\text{minfoc}\,(\gamma) = 0$*. Then* $\text{inj}(M) = 0$ *and, moreover, γ belongs to an end $E_\gamma \subset M$ of finite total curvature, $\int_{E_\gamma} |K_M| d\sigma < \infty$.*

Proof. Since γ is a geodesic ray such that $\text{minfoc}\,(\gamma) = 0$, then

$$\inf_{(p,\bar{u})\in \mathbb{S}T\gamma^\perp} e_c(p,\bar{u}) = 0 \tag{5.1}$$

with $\mathbb{S}T\gamma^\perp$ being the unitary normal bundle of γ in M. Hence, there exist a sequence of points

$$\{(p_n,\bar{u}_n)\}_{n=1}^{\infty} \subset \mathbb{S}T\gamma^\perp$$

such that

$$\lim_{n\to\infty} e_c(p_n,\bar{u}_n) = 0. \tag{5.2}$$

Since e_c is continuous, $\{(p_n,\bar{u}_n)\}_{n=1}^{\infty}$ diverges in the unitary normal bundle $\mathbb{S}T\gamma^\perp$, because on the contrary, if $(p_o,\bar{u}_0) = \lim_{n\to infinity}(p_n,\bar{u}_n)$, then $e_c(p_0,\bar{u}_0) = 0$ which is a contradiction with the fact that $exp^\perp$ is a diffeomorphism in an open set Ω_γ, (see Remark 2.4).

Taking into account that $\|\bar{u}_n\| = 1\ \forall n$, then $\{p_n\}_{n=1}^{\infty}$ diverges in M and then, as the topology of M is finite, there exists an annular end E_γ such that a divergent subsequence of $\{p_n\}_{n=1}^{\infty}$ is included in E_γ (in fact, the ray γ is included in E_γ). We are going to prove that the infimum of the injectivity radius of such a subsequence is zero and hence by applying lemma 4.1 (see remark 4.2) E_γ has finite total curvature.

Let $\alpha_{\bar{u}_n}^{p_n} : [0,\infty) \to M$ be the geodesic curve starting at p_n ($\alpha_{\bar{u}_n}^{p_n}(0) = p_n$) with initial velocity $\bar{u}_n$ ($\dot{\alpha}_{\bar{u}_n}^{p_n}(0) = \bar{u}_n$). Then, the points

$$q_n = \alpha_{\bar{u}_n}^{p_n}\left(e_c\left(p_n,\bar{u}_n\right)\right) \tag{5.3}$$

are cut-focal points of γ, where the distance from p_n is no longer minimized along $\alpha_{\bar{u}_n}^{p_n}$ after q_n. By definition, thus,

$$\mathrm{L}_n := \text{dist}_M\left(p_n,q_n\right) = e_c\left(p_n,\bar{u}_n\right). \tag{5.4}$$

Hence, by using the limit (5.2), we have that $\lim_{n\to\infty} L_n = 0$ so, for any $\epsilon > 0$, there exists N large enough such that for any $n > N$

$$\text{dist}_M\left(p_n,q_n\right) < \epsilon. \tag{5.5}$$

Therefore, there exists a divergent subsequence of cut-focal points $\{q_n\}_{n=1}^\infty$ included in the same end E_γ.

Let us consider now

$$\Omega_\gamma := \left\{(p, t\bar{u}) \in \mathbb{S}T\gamma^\perp, \quad 0 \leq t < e_c(p, \bar{u})\right\}.$$

It is known (see remark 2.4) that $\exp^\perp : \Omega_\gamma \to \exp^\perp(\Omega_\gamma)$ is a diffeomorphism and that the boundary $\partial \exp^\perp(\Omega_\gamma)$ of $\exp^\perp(\Omega_\gamma)$ is the set of cut-focal points of γ. Hence, there exists an infinite divergent sequence of $\{q_n\}_{n=1}^\infty$ in $E_\gamma \cap \partial \exp^\perp(\Omega_\gamma)$.

For these cut-focal points q_n, either there is more than one minimizing geodesic joining q_n and γ or $\ker\left(\exp^\perp_{*q}\right) \neq \{0\}$. But this last possibility implies that q_n is a focal point of γ in M, and this is not possible because γ is a totally geodesic submanifold in the negatively curved manifold M, see Proposition 2.12.

Therefore, given a point $p_n \in E_\gamma$, let us consider the cut-focal point $q_n \in E_\gamma$, being $\alpha^{p_n}_{\bar{u}_n}(t)$ the geodesic which realizes the distance $L_n = e_c(p_n, \bar{u}_n)$ from $p_n \in \gamma$ to q_n.

This geodesic meets γ orthogonally at the point p_n. Since $\exp^\perp$ is not injective when we restrict ourselves to the set of cut-focal points, then there exists another geodesic β_n minimizing the distance between q_n and γ. This geodesic will meet γ orthogonally at the point $r_n \in \gamma \subseteq E_\gamma$.

Now, let us consider the geodesic triangle $T_n \subseteq E_\gamma$ passing through the vertices p_n, q_n and r_n, formed by the union of the geodesics γ, $\alpha^{p_n}_{\bar{u}_n}$ and β_n, namely $T_n = \gamma \cup \alpha^{p_n}_{\bar{u}_n} \cup \beta_n$.

Since $\alpha^{p_n}_{\bar{u}_n}$ and β_n meet γ orthogonally, the external signed angles corresponding to the vertices p_n and r_n are $\pm\frac{\pi}{2}$ and the angle between $\alpha^{p_n}_{\bar{u}_n}$ and β_n at the point q_n, $\theta_n \in (-\pi, \pi)$. At this point, we should note that q_n is not a cusp, (namely, $|\theta_n| \neq \pi$), because in this case $\alpha^{p_n}_{u_n}$ and β_n should be the same geodesic. Therefore, the sum of the signed angles is

$$-2\pi < \text{Sum of the external signed angles of } T_n < 2\pi \tag{5.6}$$

Observe, that in the particular case when $r_n = p_n$ we have a geodesic loop.

Let us going now to see that T_n can not bound a disk. To see this, let us suppose that $\Omega \subseteq E_\gamma$ is a disk such that $\partial\Omega = T_n$. Since T_n is a piecewise curve formed by geodesic segments, by applying Gauss-Bonnet theorem

$$\int_\Omega K_M d\sigma > 2\pi\chi(\Omega) - 2\pi = 0, \tag{5.7}$$

where we have used that $\chi(\Omega) = 1$ because Ω is assumed to be a disk. But taking into account that $K_M \leq 0$, the above inequality is a contradiction.

Finally, to see that $\operatorname{inj}(M) = 0$, we are going to show that $\operatorname{inj}_M(p_n) \leq 4L_n$, and hence, $\lim\limits_{n\to\infty} \operatorname{inj}_M(p_n) \leq 4 \lim\limits_{n\to\infty} L_n = 0$. To do it, we shall prove first that $T_n \subseteq B^M_{4L_n}(p_n)$ and to see this, let us consider $x \in T_n$. Since $T_n = \gamma \cup \alpha^{p_n}_{\bar{u}_n} \cup \beta_n$, we have that $x \in \gamma$, or $x \in \alpha^{p_n}_{\bar{u}_n}$ or $x \in \beta_n$. Recall that $\operatorname{dist}_M(p_n, q_n) = \operatorname{dist}_M(q_n, r_n) = L_n = e_c(p_n, \bar{u}_n)$. Then:

If $x \in \alpha^{p_n}_{\bar{u}_n}$, then

$$\operatorname{dist}_M(p_n, x) \leq \operatorname{dist}_M(p_n, q_n) \leq L_n \tag{5.8}$$

If $x \in \beta_n$, then

$$\begin{aligned} \operatorname{dist}_M(p_n, x) \leq& \operatorname{dist}_M(p_n, q_n) + \operatorname{dist}_M(q_n, x) \\ \leq& \operatorname{dist}_M(p_n, q_n) + \operatorname{dist}_M(q_n, r_n) \\ \leq& 2L_n \end{aligned} \tag{5.9}$$

Finally, if $x \in \gamma$, then, as $\operatorname{dist}_M(r_n, p_n) \leq \operatorname{dist}_M(p_n, q_n) + \operatorname{dist}_M(q_n, r_n)$,

$$\begin{aligned} \operatorname{dist}_M(p_n, x) \leq& \operatorname{dist}_M(p_n, q_n) + \operatorname{dist}_M(q_n, r_n) + \operatorname{dist}_M(r_n, x) \\ \leq& \operatorname{dist}_M(p_n, q_n) + \operatorname{dist}_M(q_n, r_n) + \operatorname{dist}_M(r_n, p_n) \leq 4L_n \end{aligned} \tag{5.10}$$

Hence $T_n \subseteq B^M_{4L_n}(p_n)$. If $\operatorname{inj}_M(p_n) > 4L_n$, then T_n (which can not bound a disk) is included in a domain which is homeomorphic to a disk, which is a contradiction by using the Jordan-Schoenflies theorem (see [4] for instance). □

6. ON THE MINIMAL FOCAL DISTANCE OF A GEODESIC RAY

As a first corollary of Theorem 1.7 we have the following result, assuming that the surface has negative curvature. Namely:

Corollary 6.1. *Let M be a complete, non compact, and orientable surface with finite topology, non-positive Gaussian curvature $K_M \leq 0$ and such that every ray γ satisfies that* $\operatorname{minfoc}(\gamma) = 0$*. Then, M has finite total curvature, $\int_M |K_M| d\sigma < \infty$.*

Proof. The proof of Corollary 6.1 follows directly from the hypothesis that *all* the rays in M have zero focal distance and that M has finite topology. □

In this section we provide some analytic and geometric sufficient and necessary conditions for the existence of geodesic rays with zero minimal focal distance. First of all, note that as an immediate consequence of Corollary 2.15, we have the following:

Corollary 6.2. *Let M be a complete, non compact, and orientable surface with non-positive Gaussian curvature and finite area. Then, for any geodesic ray γ in M ,*

$$\operatorname{minfoc}(\gamma) = 0. \tag{6.1}$$

Hence, in the non-positively curved setting, finite area implies zero minimal focal distance for any geodesic ray. Observe, however that the converse is in general false, see for instance example 1.8 in Section §.1, where we have zero minimal focal distance for a geodesic ray and infinite area. Nevertheless, by using Theorem 1.7 we can state

Corollary 6.3. *Let M be a complete, non compact, and orientable surface with negative Gaussian curvature $K_M \leq b < 0$ and finite topological type. Given a geodesic ray γ in M,* $\operatorname{minfoc}(\gamma) = 0$ *if and only if there exist an end E_γ with finite area such that γ belongs to E_γ.*

Therefore,

Corollary 6.4. *Let M be a complete, non compact, and orientable surface with negative Gaussian curvature $K_M < 0$ and finite topological type. Suppose that every end of M has infinite area. Suppose, moreover, that a geodesic ray γ in M has* $\operatorname{minfoc}(\gamma) = 0$*. Then, there exist an end E_γ such that γ belongs to E_γ and a divergent sequence of points $\{p_i\}_{i=1}^\infty \subset E_\gamma$, such that*

$$\lim_{i\to\infty} K_M(p_i) = 0. \tag{6.2}$$

Whence, the conclusion of the above corollaries becomes clear. In a negatively curved surface of finite topological type the presence of a geodesic ray γ with zero minimal focal distance implies that there exist an end containing γ with finite area or the end is asymptotically flat (in the sense of the limit (6.2)).

Remark 6.5. From Corollaries 6.1 and 6.2 we can conclude moreover that given M a complete, non-compact and orientable surface with finite topology, finite area and non-positive curvature, then M has finite total curvature.

K. Ichihara proved in [21] that if a surface (or an end) has finite total curvature, the surface (or the end) is a parabolic surface (or end). The same is true if the surface (or end) has finite total area. Hence,

Corollary 6.6. *Let M be a negatively curved and orientable surface of finite topological type, if every end of M is an hyperbolic end, then for any geodesic ray γ in M,*

$$\operatorname{minfoc}(\gamma) > 0. \tag{6.3}$$

Moreover, if a surface (or an end) has positive fundamental tone, the surface is hyperbolic (see [20]). Recall that the fundamental tone $\lambda^*(M)$ of a surface M is defined by

$$\lambda^*(M) = \inf\{\frac{\int_M |\nabla f|^2 dA}{\int_M f^2 dA}; \, f \in L^2_{1,0}(M) \setminus \{0\}\} \tag{6.4}$$

where $L^2_{1,\,0}(M)$ is the completion of $C^\infty_0(M)$ with respect to the norm $\|\varphi\|^2 = \int_M \varphi^2 + \int_M |\nabla\varphi|^2$.

From the definition of the fundamental tone, one readily concludes that for any end $E \subset M$

$$\lambda^*(M) \leq \lambda^*(E). \tag{6.5}$$

Thus,

Corollary 6.7. *Let M be a negatively curved and orientable surface of finite topological type, if M has positive fundamental tone $\lambda^*(M) > 0$, then for any geodesic ray γ in M,*

$$\operatorname{minfoc}(\gamma) > 0. \tag{6.6}$$

REFERENCES

[1] M. T. Anderson *The compactification of a minimal submanifold by the Gauss Map.*, Preprint IEHS (1984).

[2] G. P. Bessa, L. Jorge and J. F. Montenegro *Complete submanifolds of $\mathbb{R}^n$ with finite topology.*, Communications in Analysis and Geometry, ISSN 1019-8385, Vol. 15, 4, 2007 , 725-732

[3] G. P. Bessa and M. S. Costa *On Submanifolds With Tamed Second Fundamental Form.*, Glasgow Mathematical Journal, 51, 2009, 669-680

[4] S.S. Cairns *An elementary proof of the Jordan-Schoenflies theorem*, Proc. Amer. Math. Soc. 2, (1951), 860–867.

[5] E. Calabi *Problems in Differential Geometry, (S. Kobayashi and J. Eells, Jr, eds.)*, Proc. of the United States-Japan seminarin Differential Geometry, Kyoto Japan, 1965, Nippon Hyoronsha Co., Ltd., Tokyo, (1966), 170.

[6] Huai-Dong Cao, Ying Shen, and Shunhui Zhu *The structure of stable minimal hypersurfaces in $\mathbb{R}^{n+1}$*, Mathematical Research Letters, vol 4, (1997), 637-644.

[7] M.P. Cavalcante, H. Mirandola and F. Vitorio, *The non-parabolicity of infinite volume ends*, to appear in Proc. Amer. Math. Soc., preprint arXiv:1201.6391v3, 2013.

[8] Isaac Chavel *Riemannian geometry: a modern introduction*, Cambridge Tracts in Mathematics, **108**, Cambridge University press,1993.

[9] Chen Qing. *On the volume growth and the topology of complete minimal submanifolds of a Euclidean space* J. Math. Sci. Univ. Tokyo 2 (1995), 657-669.

[10] L. Cheung, and P. Leung *The mean curvature and volume growth of compete non-compact submanifolds*, Differ. Geom. Appl. 8 (3), (1998), 251-256. 2 (1995), 657-669.

[11] T. H. Colding and W. P. Minicozzi III *The Calabi-Yau conjectures for embedded surfaces*, Annals of Mathematics, vol 167, (2008), 211-243.

[12] T. H. Colding and W. P. Minicozzi III *An excursion into Geometric Analysis*, Surv. Differ. Geom., IX, Int. Press, Somerville, M A, (2004), 83–146, math.D G /0309021.

[13] B. Coskunuzer, W. H. Meeks and G. Tinaglia *Non-properly embedded H-planes in $\mathbb{H}^3$*, Preprint , (2015), https://profmeeks.wordpress.com/

[14] M. P. do Carmo, Q. Wang and C. Xia, *Total curvature and L^2 harmonic 1-forms on complete sub manifolds in space forms*, Journal of Geometry and Physics, 60 (2010), 142-154.

[15] M. P. do Carmo, *Riemannian geometry*, Mathematics: Theory & Applications, Birkhäuser Boston Inc., Boston, MA, 1992, Translated from the second Portuguese edition by Francis Flaherty. MR 1138207 (92i:53001)

[16] H-P. Fu and H-W. Xu, *Total curvature and L^2 harmonic 1-forms on complete sub manifolds in space forms*, Geom. Dedicata 144 (2010), 129-140.

[17] A. Gray, *Tubes*, second ed., Progress in Mathematics, vol. 221, Birkhäuser Verlag, Basel, 2004, With a preface by Vicente Miquel. MR 2024928 (2004j:53001)

[18] A. Gray, *The volume of a small geodesic ball of a Riemannian manifold* , Michigan Math. J., Vol. 20, (4), (1974), 329-344.

[19] R. Greene and H. Wu *Function theory on manifolds which possess a pole.*, Lecture Notes in Math., vol. 699, Springer-Verlag, Berlin and New York, 1979.

[20] A. Grigor'yan, *Analytic and geometric background of recurrence and non-explosion of the Brownian motion on Riemannian manifolds*, Bull. Amer. Math. Soc. **36** (1999), 135–249.

[21] K. Ichihara, *Curvature, geodesics and the Brownian motion on a Riemannian manifold I; Recurrence properties*, Nagoya Math. J. **87** (1982), 101–114.

[22] L. P. Jorge and D. Koutroufiotis, *An estimate for the curvature of bounded submanifolds*, Amer. J. Math. **103** (1981), 711–725.

[23] P. Li, *Curvature and function theory on Riemannian manifolds*, Surveys in differential geometry, Surv. Differ. Geom., VII, Int. Press, Somerville, MA, 2000, pp. 375–432. MR 1919432 (2003g:53047)

[24] B. Pessoa Lima, L. Mari, J. Fabio Montenegro and F. B. Vieira, *Density and spectrum of minimal sub manifolds in space forms*, preprint arXiv:1407.5280v3, 2014.

[25] W. H. Meeks and H. Rosenberg, *The minimal lamination closure theorem*, Duke math. J., vol. 133, no. 3 (2006), 467-497.

[26] W. H. Meeks, J. Perez and a. Ros *The embedded Calabi-Yau conjectures for finite genus*, Preprint, (2015), https://profmeeks.wordpress.com/

[27] S. Muller and V. Sverak, *On surfaces of finite total curvature*, J. Differential Geometry, **42**, 2, (1995), 229-257.

[28] N. Nadirashvili *Hadamard's and Calabi's Conjectures on negatively curved and minimal surfaces*, Invent. Math., **126**, (1996), 457-465.

[29] G. DE OLIVEIRA *Compactification of minimal submanifolds of hyperbolic space.*, Comm. Analysis and Geometry, 1 (1993), 1-29.

[30] M. Rodriguez and G. Tinaglia *Non-proper Complete Minimal surfaces embedded in* $\mathbb{H}^2 \times \mathbb{R}$, International Mathematics Research Notices, rnu068, 13 pages, 2014

[31] V. Palmer *On deciding whether a submanifold is parabolic of hyperbolic using its mean curvature* , Simon Stevin Transactions on Geometry, vol 1. 131-159, Simon Stevin Institute for Geometry, Tilburg, The Netherlands, 2010.

[32] T. Sakai, *Riemannian geometry*, Translations of Mathematical Monographs, vol. 149, American Mathematical Society, Providence, RI, 1996, Translated from the 1992 Japanese original by the author. MR 1390760 (97f:53001)

[33] K. Shiohama. *Total curvatures and minimal areas of complete surfaces* Proc. Amer. Math. Soc. 94 (1985), no. 2, 310-316.

[34] M. Troyanov, *Parabolicity of Manifolds*, Siberian Advances in Mathematics **9** (1999) 125–150.

[35] J. Tysk, *Finitenes of index and total scalar curvature for minimal hypersurfaces*, Proc. Amer. Math. Soc., **105**, no. 2, (1989), 429-435.

[36] S. T. Yau, *Isoperimetric constants and the forst eigenvalue of a compact Riemannian manifold*, Ann. Sci. Ecole Norm. Sup., **8**, (1975), 487-507.

DEPARTAMENT DE MATEMÀTIQUES- IMAC, UNIVERSITAT JAUME I, CASTELLÓ, SPAIN.
E-mail address: `gimenov@uji.es`

DEPARTAMENT DE MATEMÀTIQUES- INIT, UNIVERSITAT JAUME I, CASTELLON, SPAIN.
E-mail address: `palmer@mat.uji.es`